arXiv:1111.0545v2 [math.AG] 20 Mar 2012

THE p-RANK OF THE REDUCTION $\operatorname{mod} \mathrm{p}$ OF JACOBIANS AND JACOBI SUMS

A. ÁLVAREZ*

Abstract. Let $Y_K \to X_K$ be a ramified cyclic covering of curves, where K is a cyclotomic field. In this work we study the p-rank of the reduction $\operatorname{mod} \mathrm{p}$ of a model of the jacobian of Y_K. In this way, we obtain counterparts of the Deuring polynomial, defined for elliptic curves, for genus greater than one. To carry out this study we use Jacobi sums and certain L-functions.

Contents

1. Introduction and previous notation 1
2. Cyclic extensions 4
3. L-functions 7
4. The constant term of $P_{F_{\mathfrak{p}}}^{\chi_f}(t)$ 9
5. Jacobi sums 13
6. The p-rank of jacobians 15
References 21

1. Introduction and previous notation

Previous notation: Let $K := \mathbb{Q}(\epsilon_m)$ be a cyclotomic field with ϵ_m an m-primitive root of the unity ($m > 1$), and we denote $A := \mathbb{Z}[\epsilon_m, \frac{1}{m\cdot n}]$ for some $n \in \mathbb{Z}$. We consider X_K to be a proper and geometrically irreducible curve over K, of genus g, such that it has good reduction, X, over A. Let $x_0, \cdots, x_d$ be different points of X_K given by

---

MSC: 11G30, 11G20, 11G10

* Departamento de Matemáticas. Universidad de Salamanca. Spain.

$d+1$ sections, of the structural morphism $X \to Spec(A)$, with disjoint images.

Now, let us consider $Y_K \to X_K$, a Galois ramified covering, of Galois group $G := \mathbb{Z}/m$, ramified at $x_0, \cdots, x_d$. We have that $\Sigma_{Y_K} = \Sigma_{X_K}(\sqrt[m]{f(x)})$, where Σ_{Y_K} and Σ_{X_K} are the function fields of Y_K and X_K, respectively. Let $D(f(x))$ be the principal divisor associated with $f(x) \in \Sigma_{X_K}$, $D(f(x)) = a_0 \cdot x_0 + \cdots + a_d \cdot x_d + m \cdot D$, with $0 < |a_i| < m$ and D is a divisor on X_K. Note that $a_0 + \cdots + a_d = 0 \,\mathrm{mod}\, \mathrm{m}$. Moreover, we assume that there exists a model $Y \to X$ over $Spec(A)$ for $Y_K \to X_K$.

We denote by $X_{\mathfrak{p}}$ the reduction at $\mathfrak{p} \in Spec(A)$ of X and by $Y_{\mathfrak{p}}$ a proper, smooth model for the reduction at $\mathfrak{p} \in Spec(A)$ of Y. Let us denote $k(\mathfrak{p})$ the residual field of $\mathfrak{p}$ that we assume to have $q = p^h$ elements, with $(p, m) = 1$. We also assume that $Y_{\mathfrak{p}} \to X_{\mathfrak{p}}$ is a ramified Galois covering of group $\mathbb{Z}/m$, ramified at $\overline{x}_0, \cdots, \overline{x}_d$, the reduction of $x_0, \cdots, x_d$ at $\mathfrak{p}$ and that $\Sigma_{Y_{\mathfrak{p}}} = \Sigma_{X_{\mathfrak{p}}}(\sqrt[m]{f(x)})$ where $D(f(x)) = a_0 \cdot \overline{x}_0 + \cdots + a_d \cdot \overline{x}_d + m \cdot \overline{D}$, with $\overline{D}$ a divisor on $X_{\mathfrak{p}}$. Here, we denote by $\Sigma_{Y_{\mathfrak{p}}}$, $\Sigma_{X_{\mathfrak{p}}}$ the function fields of $Y_{\mathfrak{p}}$ and $X_{\mathfrak{p}}$, respectively.

We study the characteristic polynomials of the p^h-Frobenius morphism, $F_{\mathfrak{p}}$, of a $\mathbb{Q}_l[\epsilon_m]$-module associated with the l-adic cohomology group $H^1(\overline{Y}_{\mathfrak{p}}, \mathbb{Q}_l)^\vee := Hom(\mathbb{Q}_l/\mathbb{Z}_l, Pic^0_{Y_{\mathfrak{p}}}(\mathbb{F})) \otimes_{\mathbb{Z}_l} \mathbb{Q}_l$. These polynomials are essentially given by certain incomplete L-functions of $X_{\mathfrak{p}}$, $\overline{Y}_{\mathfrak{p}} = Y_{\mathfrak{p}} \otimes_{\mathbb{F}_{\mathfrak{p}}} \mathbb{F}$.

The constant term of these polynomials is given by jacobi sums. This is proved in [D], 5 and Proposition 10.12.1, because these terms are the local constants of an Dirichlet L-series, defined over $\Sigma_{X_{\mathfrak{p}}}$ and for a character χ_f of G. In this work, we make explicit calculations of these terms by means of geometric methods; generalized jacobians, Riemann-Roch Theorem...

By using properties of Jacobi sums we prove that if m is a prime integer and $t \in \{1, \cdots, m-1\}$ with

$$[< \frac{a_1}{m} > + \cdots + < \frac{a_d}{m} >] \neq [< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >],$$

then the jacobian of $Y_{\mathfrak{p}}$ is not a supersingular abelian variety when $m | p-1$. $[\frac{a}{b}]$ denotes the integer part of the fraction $\frac{a}{b}$ and $< \frac{a}{b} > := \frac{a}{b} - [\frac{a}{b}]$. We say that an abelian variety A over $F_{\mathfrak{p}}$ is supersingular if the eigenvalues of the Frobenius morphism $F_{\mathfrak{p}}$ are $\zeta \cdot q^{1/2}$, ζ being a root of the unity.

We also prove that if m is a prime integer and $t \in \{1, \cdots, m-1\}$ with

$$[< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >] \neq 0,$$

$X_{\mathfrak{p}} = \mathbb{P}_1$ and hence $Y_{\mathfrak{p}}$ does not have p-rank 0.

Where, we say that $Y_{\mathfrak{p}}$ has p-rank 0 if $\mathrm{Ker[p]_{red}} = \mathrm{Spec(k(\mathfrak{p}))}$, $[p]$ being the multiplication by p on $\mathrm{Pic}^0_{\mathrm{Y}_{\mathfrak{p}}}$.

When m is a prime integer and p inert in $\mathbb{Z}[\epsilon_m]$, we prove that $Y_{\mathfrak{p}}$ has p-rank 0 if and only if $X_{\mathfrak{p}}$ has p-rank 0 and the coefficients, up to the constant term, of the above characteristic polynomials are 0 modulo p. As a corollary, we deduce that the proper, smooth model of the curve

$$y^m - x^{a_1}(x-1)^{a_2}(x-\alpha_1)^{a_3} \cdots (x - a_{d-2})^{a_d},$$

defined on $k(\mathfrak{p}) = \mathbb{F}_{p^{m-1}}$, has p-rank 0 if and only if $\alpha_1, \cdots, \alpha_{d-2}$ satisfy a system of $d-2$ algebraic equations defined on $\mathbb{F}_{p^{m-1}}$

$$p_1(\alpha_1, \cdots, \alpha_{d-2}) = 0, \cdots, p_{d-2}(\alpha_1, \cdots, \alpha_{d-2}) = 0.$$

Here, $0 < |a_i| < m$ for each $1 \leq i \leq d-2$, and $a_1 + \cdots + a_d \neq 0 \bmod m$. When $d = 3$, $a_1 = a_2 = a_3 = 1$ and $m = 2$, one obtains the Deuring polynomial, defined for the elliptic curve $y^2 - x(x-1)(x-\lambda)$,

$$H(\lambda) = (-1)^n \sum_{i=0}^{n} \binom{n}{i}^2 \lambda^i, (\text{with} \quad n = p - 1/2).$$

We also study when $Y_{\mathfrak{p}}$ has p-rank 0 in the case that p splits completely in $\mathbb{Z}[\epsilon_m]$.

For Fermat curves, (with our notation $g = 0$, $d = 2$), in [A2] the torsion on jacobians is studied in terms of theta divisors, in [N] the supersingularity of these curves is studied, and the article [U] is devoted, by means of jacobi sums, to study the rank of jacobians of Fermat curves over function fields. In [L] is studied the p-torsion of abelian varieties in characteristic p. In the article [NG] is proved that the set of ordinary abelian varieties is dense in the special fiber, modulo p, of certain moduli space of principally polarized abelian varieties.

We now give some general notation that we shall use along this work. Let s be a global section of a line bundle L on $X_{\mathfrak{p}}$. We also denote by s the morphism, of $\mathcal{O}_{X_{\mathfrak{p}}}$-modules, $\mathcal{O}_{X_{\mathfrak{p}}} \to L$, such that $1 \to s$. Here, $\mathcal{O}_{X_{\mathfrak{p}}}$ denotes the sheaf of rings associated with $X_{\mathfrak{p}}$.

We denote by E and $\mathfrak{m}$ the effective divisor $\overline{x}_0 + \cdots + \overline{x}_d$ on $X_{\mathfrak{p}}$ and the ideal associated with E inside $\mathcal{O}_{X_{\mathfrak{p}}}$, respectively, and $T := \{\overline{x}_0, \cdots, \overline{x}_d\}$.

If $z \in k(\mathfrak{p})^{\times} = \mathbb{F}_{p^h}^{\times}$, then we denote by $\chi_{\mathfrak{p}}(z)$ the unique m-root of the unity such that $\chi_{\mathfrak{p}}(z) = z^{\frac{p^h-1}{m}} \bmod \mathfrak{p}$.

2. CYCLIC EXTENSIONS

Let $\mathrm{Pic}^0_{X_{\mathfrak{p}},\mathfrak{m}}$ be the generalized jacobian for $\mathfrak{m}$. $\mathrm{Pic}^0_{X_{\mathfrak{p}},\mathfrak{m}}$ is a scheme over $k(\mathfrak{p})$ that represents isomorphism classes of pairs $(L, \iota_{\mathfrak{m}})$ ($\mathfrak{m}$-level structures), where L is a line bundle over $X_{\mathfrak{p}}$ and $\iota_{\mathfrak{m}} : L \to \mathcal{O}_{X_{\mathfrak{p}}}/\mathfrak{m}$ is a surjective morphism of $\mathcal{O}_{X_{\mathfrak{p}}}$-modules, we say that two level structures $(L, \iota_{\mathfrak{m}})$ and $(L', \iota'_{\mathfrak{m}})$ are equivalents when there exists an isomorphism of line bundles $u : L \to L'$ such that $\iota'_{\mathfrak{m}} \cdot u = \iota_{\mathfrak{m}}$. The equivalence classes of level structures are in one-to-one correspondence with the $\mathfrak{m}$-equivalence classes of divisors on $X_{\mathfrak{p}}$ supported outside T; two divisors

D and D', are $\mathfrak{m}$-equivalent if there exists a $g \in \Sigma_{X_\mathfrak{p}}^{\mathfrak{m}}$ with $D - D' = D(g)$ where $\Sigma_{X_\mathfrak{p}}^{\mathfrak{m}} := \{g \in \Sigma_{X_\mathfrak{p}} \text{ such that } g \equiv 1 \bmod \mathfrak{m}\}$.

We call the (projective) space of $\mathfrak{m}$-sections of a level structure $(L, \iota_\mathfrak{m})$, and we denote by $H^0_\mathfrak{m}((L, \iota_\mathfrak{m}))$, to the space of sections $s : \mathcal{O}_{X_K} \to L$ such that $\iota_\mathfrak{m} \cdot s = \pi$, $\pi : \mathcal{O}_{X_\mathfrak{p}} \to \mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m}$ being the natural epimorphism. The effective $\mathfrak{m}$-equivalent divisors associated with $(L, \iota_\mathfrak{m})$ is given by the zero locus of the $\mathfrak{m}$-sections of $(L, \iota_\mathfrak{m})$. Moreover, if $s \in H^0_\mathfrak{m}((L, \iota_\mathfrak{m}))$ then $H^0_\mathfrak{m}((L, \iota_\mathfrak{m})) = s + H^0(X_\mathfrak{p}, L(-E))$.

We denote by I_T, I_T^0 and $O_T^\times$ the ideles, ideles of degree 0 and integer ideles on $\Sigma_{X_\mathfrak{p}}$ outside T, respectively.

According class field theory, the cyclic extension $\Sigma_{Y_\mathfrak{p}}/\Sigma_{X_\mathfrak{p}}$ of Galois group $\mathbb{Z}/m$, where $\Sigma_{Y_\mathfrak{p}} = \Sigma_{X_\mathfrak{p}}(\sqrt[m]{f(x)})$, gives an epimorphism of groups

$$\frac{I_T^0}{(\Sigma_{X_\mathfrak{p}}^{\mathfrak{m}})^\times \cdot O_T^\times} \to \mathbb{Z}/m.$$

This morphism is given by the Artin map

$$(\quad, \Sigma_{Y_\mathfrak{p}}/\Sigma_{X_\mathfrak{p}}) : \frac{I_T}{(\Sigma_{X_\mathfrak{p}}^{\mathfrak{m}})^\times \cdot O_T^\times} \to \mathbb{Z}/m.$$

In this way there exists a divisor D_1 of degree 1 on $X_\mathfrak{p}$ with

$$(D_1, \Sigma_{Y_\mathfrak{p}}/\Sigma_{X_\mathfrak{p}}) = 1.$$

We consider the p^h-Frobenius morphism ($p^h = \#k(\mathfrak{p})$), $F_\mathfrak{p}$, and the Lang isogeny, $P := F_\mathfrak{p} - Id : \mathrm{Pic}^0_{X_\mathfrak{p},\mathfrak{m}} \to \mathrm{Pic}^0_{X_\mathfrak{p},\mathfrak{m}}$. Bearing in mind the divisor of degree 1, D_1, we have an immersion $X_\mathfrak{p} \setminus T \to \mathrm{Pic}^0_{X_\mathfrak{p},\mathfrak{m}}$. By class field theory for function fields over finite fields, (c.f. [S]), $P^{-1}(X_\mathfrak{p} \setminus T)$ gives the $\mathfrak{m}$-ray class field $H_\mathfrak{m}$ for $\Sigma_{X_\mathfrak{p}}$. It is a Galois extension of Galois group isomorphic to $\mathrm{Pic}^0_{X_\mathfrak{p},\mathfrak{m}}(k(\mathfrak{p})) = \frac{I_T^0}{(\Sigma_{X_\mathfrak{p}}^{\mathfrak{m}})^\times \cdot O_T^\times}$. This group is the group of the $\mathfrak{m}$-equivalence classes of divisors on $X_\mathfrak{p}$ supported outside T.

The morphism of forgetting the level structure, $(L, \iota_{\mathfrak{m}}) \to L$, gives the exact sequence of schemes in groups

$$1 \to (\mathbb{G}_m) \times \overset{d+1}{\overset{\smile}{\cdots}} \times \mathbb{G}_m / \mathbb{G}_m \to \mathrm{Pic}^0_{X_{\mathfrak{p}},\mathfrak{m}} \to \mathrm{Pic}^0_{X_{\mathfrak{p}}} \to 1$$

and thus we have the exact sequence of groups,

$$1 \to k(\overline{x}_0)^\times \times \cdots \times k(\overline{x}_d))^\times / k(\mathfrak{p})^\times \xrightarrow{\eta} \frac{I^0_T}{(\Sigma^{\mathfrak{m}}_{X_{\mathfrak{p}}})^\times \cdot O^\times_T} \to \frac{I^0_T}{\Sigma^\times_{X_{\mathfrak{p}}} \cdot O^\times_T} \to 1.$$

Note that the residual fields $k(\overline{x}_i)$ are isomorphic to $k(\mathfrak{p})$. We have

$$\chi_f(\eta(z_0, \cdots, z_d), \Sigma_{Y_{\mathfrak{p}}} / \Sigma_{X_{\mathfrak{p}}}) = \chi_{\mathfrak{p}}^{-a_0}(z_0) \cdots \chi_{\mathfrak{p}}^{-a_d}(z_d),$$

where, if $\sigma \in \frac{I^0_T}{(\Sigma_{X_{\mathfrak{p}}})^\times \cdot O^\times_T}$ and $\sigma(\sqrt[m]{f(x)}) = z \cdot (\sqrt[m]{f(x)})$ (via the Artin symbol), with $z \in k(\mathfrak{p})^\times = \mathbb{F}^\times_{p^h}$, then we define $\chi_f(\sigma) := \chi_{\mathfrak{p}}(z)$. Recall that $\chi_{\mathfrak{p}}(z)$ is the unique m-root of the unity satifying $\chi_{\mathfrak{p}}(z) = z^{q^h - 1/m} \mod \mathfrak{p}$ and that $D(f(x)) = a_0 \cdot \overline{x}_0 + \cdots + a_d \cdot \overline{x}_d + m \cdot \overline{D}$, with $0 < |a_i| < m$ and $\overline{D}$ is a divisor on $X_{\mathfrak{p}}$.

We have an isomorphism

$$k(\overline{x}_0)^\times \times \cdots \times k(\overline{x}_d))^\times / k(\mathfrak{p})^\times \simeq k(\overline{x}_1)^\times \times \cdots \times k(\overline{x}_d))^\times$$

and thus we can assume that $z_0 = 1$.

Remark 2.1. Bearing in mind the notation of the introduction, if $Y_K \to X_K$ is a Galois ramified covering, of Galois group $G := \mathbb{Z}/m$, ramified at $x_0, \cdots, x_d$, it is not difficult to show that there exists $A := \mathbb{Z}[\epsilon_m, \frac{1}{m \cdot n}]$ for some $n \in \mathbb{Z}$, such that $Y_K \to X_K$ has a model, $Y \to X$ over $Spec(A)$. Moreover, $x_0, \cdots, x_d$ are given by $d+1$-sections $Spec(A) \to X$, with a disjoint image, and for each $\mathfrak{p} \in Spec(A)$, $Y_{\mathfrak{p}} \to X_{\mathfrak{p}}$ is a Galois ramified covering of group $\mathbb{Z}/m$ ramified over $\overline{x}_0, \cdots, \overline{x}_d$, the reduction of $x_0, \cdots, x_d$ at $\mathfrak{p}$. To obtain $Y \to X$, it suffices to consider a model over A of the generalized jacobian of X_K for the divisor $x_0 + \cdots + x_d$.

3. L-functions

This section is devoted to studying certain incomplete L-functions of the curve $X_{\mathfrak{p}}$ over $k(\mathfrak{p})$ for each $\mathfrak{p} \in Spec(A)$. We follow [A1] and [T]. We shall calculate the characteristic polynomial of $F_{\mathfrak{p}}$ as an endomorphism of a certain $\mathbb{Q}_l[\epsilon_m]$-module obtained from the $\mathbb{Z}/m$-module $H^1(\overline{Y}_{\mathfrak{p}}, \mathbb{Q}_l)^{\vee}$.

We now consider the T-incomplete L-function

$$\theta_{H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}},T}(t) = \prod_{x \in |X_{\mathfrak{p}}| \setminus T} (1 - F_x \cdot t^{deg(x)})^{-1},$$

where $F_x \in G_{\mathfrak{m}} := \frac{I_T^0}{(\Sigma_{X_{\mathfrak{p}}}^{\mathfrak{m}})^{\times} \cdot O_T^{\times}}$ is the Frobenius element, for the point $x \in |X_{\mathfrak{p}}| \setminus T$, considered as an element of the Galois group, $G_{\mathfrak{m}}$, of the ray class field for $\mathfrak{m}$, $H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}}$. We have that $F_x = (t_x, H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}})$, where $(\quad, H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}})$ is the Artin symbol for the Galois extension $H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}}$, and t_x is a local parameter for x.

We denote by σ_N the element of the Galois group of the extension $H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}}$ associated with N via the Artin symbol, N being a divisor supported outside T and with class $[N] \in G_{\mathfrak{m}}$. Similar to [A1] 4.1.1, we can compute this L-function in terms of $G_{\mathfrak{m}}$:

$$\theta_{H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}},T}(t) = \sum_{[N] \in G_{\mathfrak{m}}} \sigma_N \cdot \Big(\sum_{i=0}^{2g+d-1} L_{\mathfrak{m}}(N,i) \cdot t^i + \sum_{j \geq 0} q^{g+j} \cdot t^{2g+d+j} \Big) =$$

$$= \sum_{i=0}^{2g+d-1} \sum_{[N] \in G_{\mathfrak{m}}} (L_{\mathfrak{m}}(N,i) \cdot \sigma_N) \cdot t^i + \Big(\sum_{[N] \in G_{\mathfrak{m}}} \sigma_N \Big) \Big(\frac{q^g \cdot t^{2g+d}}{1 - qt} \Big) =$$

$$= \sum_{i=0}^{2g+d-1} \sum_{D, \mathrm{deg(D)=i}} (D, H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}}) \cdot t^{2g+d-1-i} + \Big(\sum_{[N] \in G_{\mathfrak{m}}} \sigma_N \Big) \Big(\frac{q^g \cdot t^{2g+d}}{1 - qt} \Big),$$

where D denotes effective divisors on $X_{\mathfrak{p}}$ with support outside T, $L_{\mathfrak{m}}(N,i)$ denotes the cardinal of the set of effective divisors on $X_{\mathfrak{p}}$ supported outside T and $\mathfrak{m}$-equivalents to $N + i \cdot D_1$. If $D = n_1 \cdot y_1 +$

$\cdots + n_r \cdot y_r$ is a divisor on $X_{\mathfrak{p}}$ with support outside T, we define

$$(D, H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}}) := \prod_{i=1}^{r} (t_{y_i}, H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}})^{n_i}$$

where t_{y_i} is a local parameter for y_i.

Note that if $(L, \iota_{\mathfrak{m}})$ is a level structure associated with $N + i \cdot D_1$ then $L_{\mathfrak{m}}(N, i) = \# H^0_{\mathfrak{m}}((L, \iota_{\mathfrak{m}}))$. Thus either $L_{\mathfrak{m}}(N, i) = 0$ or $L_{\mathfrak{m}}(N, i) = \# H^0(X_{\mathfrak{p}}, N + i \cdot D_1 - E)$. The divisor D_1 is defined in section 2.

For ease in the notation, we assume that m and l are integer primes, such that l is inert within $\mathbb{Z}[\epsilon_m]$. Thus, $\mathbb{Q}_l[\epsilon_m]$ is a field and $\mathbb{Q}_l[G] = \mathbb{Q}_l \times \mathbb{Q}_l[\epsilon_m]$, $(G := \mathbb{Z}/m)$. Let us denote $e_1 \in \mathbb{Z}[G]$, the idempotent element, such that $e_1 \cdot \mathbb{Q}_l[G] = 0 \times \mathbb{Q}_l[\epsilon_m]$. Let us consider the first l-adic cohomology group $H^1(Y_{\mathfrak{p}}, \mathbb{Q}_l)^{\vee}$. This is a $\mathbb{Q}_l[G]$-module and $e_1 \cdot H^1(Y_{\mathfrak{p}}, \mathbb{Q}_l)^{\vee}$ is a $\mathbb{Q}_l[\epsilon_m]$-module.

The p^h-Frobenius morphism $F_{\mathfrak{p}}$ is an endomorphism of this $\mathbb{Q}_l[\epsilon_m]$-module. By [T] 3.5, since $Y_{\mathfrak{p}} \to X_{\mathfrak{p}}$ is ramified on $\overline{x}_0, \cdots, \overline{x}_d$ we have that the characteristic polynomial of $F_{\mathfrak{p}}$, $P^{\chi_f}_{F_{\mathfrak{p}}}(t) := det(F_{\mathfrak{p}} - t) \in \mathbb{Z}[\epsilon_m][t]$, is the polynomial

$$(*) \chi_f(t^{2g+d-1} \cdot \theta_{H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}}, T}(\frac{1}{t})) = \sum_{i=0}^{2g+d-1} \sum_{[N] \in G_{\mathfrak{m}}} L_{\mathfrak{m}}(N, i) \cdot \chi_f(\sigma_N) \cdot t^{2g+d-1-i}$$

$$= \sum_{i=0}^{2g+d-1} \sum_{D, \mathrm{deg(D)=i}} \chi_{\mathfrak{p}}(f(D)) t^{2g+d-1-i},$$

where χ_f is defined in section 2 and D denotes effective divisors on $X_{\mathfrak{p}}$ with support outside T. If $D = n_1 \cdot y_1 + \cdots + n_r \cdot y_r$ is a divisor on $X_{\mathfrak{p}}$ then we define

$$f(D) := \prod_{i=1}^{r} f(y_i)^{n_i \cdot \frac{p^{h \cdot \mathrm{deg(y_i)}} - 1}{p^h - 1}}.$$

If $x \in \mathrm{sup(D(f))} \setminus \mathrm{T}$ then we define $f(x)$ by considering a divisor D' linearly $\mathfrak{m}$-equivalent to x and $\mathrm{sup(D')} \cap \mathrm{sup(D(f))} = \emptyset$.

The characteristic polynomial of $F_{\mathfrak{p}}$, where $F_{\mathfrak{p}}$ is considered as an endomorphism of the $\mathbb{Q}_l$-vector space $H^1(\overline{Y}_{\mathfrak{p}}, \mathbb{Q}_l)^{\vee}$, is given by

$$p_0(t) \cdot \prod_{1 \leq j < m} P_F^{\chi_f^j}(t) \in \mathbb{Z}[t].$$

Here, $p_0(t)$ denotes the characteristic polynomial of $F_{\mathfrak{p}}$ as an endomorphism of the $\mathbb{Q}_l$-vector space $H^1(\overline{X}_{\mathfrak{p}}, \mathbb{Q}_l)^{\vee}$.

4. The constant term of $P_{F_{\mathfrak{p}}}^{\chi_f}(t)$

In this section we shall calculate the constant term of the polynomial $P_{F_{\mathfrak{p}}}^{\chi_f}(t)$. This means, by (*), to calculating $\sum_{[N]\in G_{\mathfrak{m}}} L_{\mathfrak{m}}(N, 2g+d-1) \cdot \chi_f(\sigma_N)$. We shall calculate $L_{\mathfrak{m}}(N, 2g+d-1)$ for each $[N] \in G_{\mathfrak{m}}$. Let κ be a divisor of degree $2g-2$ associated with the canonical sheaf of $X_{\mathfrak{p}}$. We have two cases for the cardinal

$$\# H^0(X_{\mathfrak{p}}, N + (2g+d-1)\cdot D_1 - E).$$

It is $= q^g$ in the case that $N + (2g+d-1)\cdot D_1 - E$ is linearly equivalent to κ and $= q^{g-1}$ in the other case.

Lemma 4.1. *Let* $(L, \iota_{\mathfrak{m}})$ *be a level structure where* $\deg(\mathrm{L}) = 2\mathrm{g}+\mathrm{d}-1$. *If* L *is a line bundle not isomorphic to* $\mathcal{O}_{X_{\mathfrak{p}}}(\kappa+E)$ *then* $\# H^0_{\mathfrak{m}}((L, \iota_{\mathfrak{m}})) = q^{(g-1)}$.

Proof. By taking global sections on the exact sequence

$$0 \to L(-E) \to L \overset{\iota_{\mathfrak{m}}}{\to} \mathcal{O}_{X_{\mathfrak{p}}}/\mathfrak{m} \to 0$$

we obtain the exact sequence of vector spaces

$$0 \to H^0(X_{\mathfrak{p}}, L(-E)) \to H^0(X_{\mathfrak{p}}, L) \overset{H^0(\iota_{\mathfrak{m}})}{\to} \mathcal{O}_{\mathfrak{m}} \to 0,$$

where we have denoted $\mathcal{O}_{\mathfrak{m}} := H^0(X_{\mathfrak{p}}, \mathcal{O}_{X_{\mathfrak{p}}}/\mathfrak{m})$. Thus, we have an isomorphism of $\mathbb{F}_{p^h}$-vector spaces

$$H^0(\iota_{\mathfrak{m}}) : H^0(X_{\mathfrak{p}}, L)/H^0(X_{\mathfrak{p}}, L(-E)) \to \mathcal{O}_{\mathfrak{m}}.$$

Therefore, there exists a section $s : \mathcal{O}_{X_\mathfrak{p}} \to L$ such that $H^0(\iota_\mathfrak{m})(s) = \pi(1)$; thus we have that $H^0_\mathfrak{m}(L, \iota_\mathfrak{m}) \neq 0$ and this cardinal is p^{g-1}. Here, π is the natural epimorphism $\mathcal{O}_{X_\mathfrak{p}} \to \mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m}$. $\square$

We now study the case when $L \simeq \mathcal{O}_{X_\mathfrak{p}}(\kappa + E)$.

We denote by $\pi_{\overline{x}_0}$ the surjective morphism of modules $\mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m} \to \mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m}_{\overline{x}_0}$, with $\mathfrak{m}_{\overline{x}_0}$ the maximal ideal associated with $\overline{x}_0$. We denote $\iota_{\overline{x}_0} := \pi_{\overline{x}_0} \cdot \iota_\mathfrak{m}$. Recall that $E = \overline{x}_0 + \cdots + \overline{x}_d$ and $\mathfrak{m}$ is the ideal within $\mathcal{O}_{X_\mathfrak{p}}$ associated with E.

We shall study when a level structure $(L, \iota_\mathfrak{m})$, with L a line bundle isomorphic to $\mathcal{O}_{X_\mathfrak{p}}(\kappa + E)$, has an $\mathfrak{m}$-section.

Bearing in mind that if $\lambda \in \mathbb{F}_\mathfrak{p}^\times$ then $(L, \iota_\mathfrak{m})$ and $(L, \lambda \cdot \iota_\mathfrak{m})$ are isomorphic level structures, we can fix the morphism $\iota_{\overline{x}_0} := \pi_{\overline{x}_0} \cdot \iota_\mathfrak{m} : L \to \mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m} \to \mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m}_{\overline{x}_0}$: For the level structures $(\mathcal{O}_{X_\mathfrak{p}}, \iota_\mathfrak{m})$, we fix $\iota_{\overline{x}_0}(1) = 1$. Let us consider $\overline{D}$ an effective divisor with $\overline{x}_0 \notin \mathrm{supp}(\overline{\mathrm{D}})$, to fix the $\overline{x}_0$-level structures for $\mathcal{O}_{X_\mathfrak{p}}(\overline{D})$ it suffices to consider the already fixed $\mathfrak{m}_{\overline{x}_0}$-level structure for $\mathcal{O}_{X_\mathfrak{p}}$ and the natural inclusion $\mathcal{O}_{X_\mathfrak{p}} \hookrightarrow \mathcal{O}_{X_\mathfrak{p}}(\overline{D})$. If L is of degree $2g + d - 1$, we can obtain an effective divisor, $\overline{D}$, with support outside $\overline{x}_0$, such that L is isomorphic to $\mathcal{O}_{X_\mathfrak{p}}(\overline{D})$. We consider a $\overline{D}$ linearly equivalent to $\kappa + E$, with $\overline{x}_0 \notin \sup(\overline{D})$, and we fix $\iota_{\overline{x}_0}$ for $L = \mathcal{O}_{X_\mathfrak{p}}(\overline{D})$.

We denote by E' the divisor $\overline{x}_1 + \cdots + \overline{x}_d$, by $\mathfrak{m}'$ the ideal associated with E', and $\mathcal{O}_{\mathfrak{m}'} := H^0(X_\mathfrak{p}, \mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m}')$.

Because $\mathcal{O}_\mathfrak{m} = k(\overline{x}_0) \times \mathcal{O}_{\mathfrak{m}'}$, for each $\mathfrak{m}$-level structure $(L, \iota_\mathfrak{m})$ we have that $\iota_\mathfrak{m} = \iota_{\overline{x}_0} \times \iota_{\mathfrak{m}'}$, recall that $\iota_{\overline{x}_0}$ is fixed and that $L \simeq \mathcal{O}_{X_\mathfrak{p}}(\kappa + E)$.

Note that $H^0(X_\mathfrak{p}, L(-E)) = H^0(X_\mathfrak{p}, L(-E'))$ thus via $H^0(\iota_{\mathfrak{m}'})$ we have the isomorphism of vector spaces

$$H^0(X_\mathfrak{p}, L)/H^0(X_\mathfrak{p}, L(-E)) = H^0(X_\mathfrak{p}, L)/H^0(X_\mathfrak{p}, L(-E')) \overset{H^0(\iota_{\mathfrak{m}'})}{\to} \mathcal{O}_{\mathfrak{m}'},$$

we denote this isomorphism by $[\iota_{\mathfrak{m}'}]$.

Let $H^0(\iota_{\overline{x}_0})$ be the linear morphism $H^0(X_\mathfrak{p}, L) \overset{H^0(\iota_\mathfrak{m})}{\to} \mathcal{O}_\mathfrak{m} \overset{\pi_{\overline{x}_0}}{\to} k(\overline{x}_0)$, we now choose an $\mathfrak{m}'$-level structure for L, (L, ι'). We consider the $\mathbb{F}_{p^h}$-linear form

$$\omega := H^0(\iota_{\overline{x}_0}) \cdot [\iota']^{-1} : \mathcal{O}_{\mathfrak{m}'} \simeq H^0(X_\mathfrak{p}, L)/H^0(X_\mathfrak{p}, L(-E)) \to k(\overline{x}_0) = \mathbb{F}_{p^h}.$$

Here, we must bear in mind that $H^0(\iota_{\overline{x}_0})(H^0(X_\mathfrak{p}, L(-E))) = 0$.

By considering the standard basis for the $\mathbb{F}_{p^h}$-vector space $\mathcal{O}_{\mathfrak{m}'} = k(\overline{x}_1) \times \cdots \times k(\overline{x}_d)$ we have that $\omega(z_1, \cdots, z_d) = \lambda_1 \cdot z_1 + \cdots + \lambda_d \cdot z_d$. Moreover, since $\mathrm{deg(L) = 2g + d - 1}$ and

$$\mathrm{Ker}(\omega) = [\iota'](\mathrm{H^0(X_\mathfrak{p}, L(-\overline{x}_0)/H^0(X_\mathfrak{p}, L(-E)))})$$

we have that for $1 \leq i \leq d$

$$H^0(X_\mathfrak{p}, L(-\overline{x}_0)/H^0(X_\mathfrak{p}, L(-E)) \neq H^0(X_\mathfrak{p}, L(-\overline{x}_0 - \overline{x}_i)/H^0(X_\mathfrak{p}, L(-E)),$$

thus, $\lambda_1, \cdots, \lambda_d \neq 0$. Accordingly, by changing the choice of the $\mathfrak{m}'$-level structure ι', we can assume that $\lambda_1 = \cdots = \lambda_d = -1$.

Note that if $(L, \iota_\mathfrak{m})$ is an $\mathfrak{m}$-level structure, then there exists $z \in \mathcal{O}^\times_{\mathfrak{m}'}$ with $\iota_\mathfrak{m} = \iota_{\overline{x}_0} \times z \cdot \iota'$. Recall that here $L \simeq \mathcal{O}_{X_\mathfrak{p}}(\kappa + E)$.

Lemma 4.2. *By using the above notations,* $H^0_\mathfrak{m}((L, \iota_{\overline{x}_0} \times z \cdot \iota')) \neq 0$ *if and only if* $\omega(z^{-1}) = 1$.

Proof. If there exists a section s of L such that the diagram

$$\begin{tikzcd}
\mathcal{O}_{X_\mathfrak{p}} \arrow[r, "s"] \arrow[dr, "\pi'"] & L \arrow[d, "z\cdot\iota'"] \\
& \mathcal{O}_{X_\mathfrak{p}}/\mathfrak{m}'
\end{tikzcd}$$

is commutative, π' being the natural epimorphism, we have that the class of the section s in $H^0(X_\mathfrak{p}, L)/H^0(X_\mathfrak{p}, L(-E'))$ is $[z \cdot \iota']^{-1}(1)$. Recall that $[z \cdot \iota']$ is the isomorphism, given by

$$H^0(X_\mathfrak{p}, L)/H^0(X_\mathfrak{p}, L(-E)) = H^0(X_\mathfrak{p}, L)/H^0(X_\mathfrak{p}, L(-E')) \overset{H^0(z\cdot\iota')}{\to} \mathcal{O}_{\mathfrak{m}'}.$$

Moreover, as the diagram

$$\begin{tikzcd}
\mathcal{O}_{X_{\mathfrak{p}}} \arrow[r, "s"] \arrow[dr, "\pi_{\overline{x}_0}"] & L \arrow[d, "\iota_{\overline{x}_0}"] \\
 & k(\overline{x}_0)
\end{tikzcd}$$

must also be commutative, we have that $H^0(\iota_{\overline{x}_0})(s) = 1$, recall that $\iota_{\overline{x}_0}$ is alrady fixed. Thus, we deduce that $H^0(\iota_{\overline{x}_0})([z \cdot \iota']^{-1}(1)) = 1$. Therefore,

$$1 = H^0(\iota_{\overline{x}_0})([z \cdot \iota']^{-1}(1)) = H^0(\iota_{\overline{x}_0})([\iota']^{-1}(z^{-1})) = \omega(z^{-1}).$$

Reciprocally, it suffices to consider a section s of L in the class of $[z \cdot \iota']^{-1}(1) \in H^0(X_{\mathfrak{p}}, L)/H^0(X_{\mathfrak{p}}, L(-E'))$. □

We denote by M a divisor of degree 0 with support outside T such that the $\mathfrak{m}$-level structure associated with $M + (2g + d - 1) \cdot D_1$ is $(\mathcal{O}_{X_{\mathfrak{p}}}(\kappa + E), \iota_{x_0} \times \iota')$, and we recall that σ_M is the element of the Galois group of the extension $H_{\mathfrak{m}}/\Sigma_{X_{\mathfrak{p}}}$ given, via the Artin symbol, by the class $[M] \in \frac{I_T^0}{(\Sigma_{X_{\mathfrak{p}}}^{\mathfrak{m}})^\times \cdot O_T^\times}$.

From Lemmas 4.1 and 4.2 we deduce, [D]:

Lemma 4.3. *The term constant of the characteristic polynomial $P_{F_{\mathfrak{p}}}^{\chi_f}(t)$, defined in section 2, is*

$$\sum_{[N] \in G_{\mathfrak{m}}} L_{\mathfrak{m}}(N, 2g + d - 1) \cdot \chi_f(\sigma_N) = \sum_{z \in \mathcal{O}_{\mathfrak{m}'}^\times,\, \omega(z) = 1} q^g \cdot \chi_f(\sigma_{z^{-1} \cdot M}) +$$

$$+ \sum_{\substack{[N] \in G_{\mathfrak{m}} \\ \text{with } N + (2g+d-1)D_1 \nsim \kappa + E}} q^{g-1} \cdot \chi_f(\sigma_N).$$

We have denoted $z^{-1} \cdot M$ instead of $\eta(z^{-1}) \cdot [M]$ and we have beared in mind that the set of classes of level structures for the line bundle

$$\mathcal{O}_{X_{\mathfrak{p}}}(\kappa + E - (2g + d - 1)D_1)$$

is given by $\{\eta(z^{-1}) \cdot [M]\}_{z \in \mathcal{O}_{\mathfrak{m}'}^\times} \subset \frac{I_T^0}{(\Sigma_{X_{\mathfrak{p}}}^{\mathfrak{m}})^\times \cdot O_T^\times}$.

5. Jacobi sums

In the first part of this section we follow, [W]. Let p be an integer prime, and $\mathfrak{p}$ a prime on $\mathbb{Z}[\epsilon_m]$ above p with residual field $\mathbb{F}_{p^h}$, a finite field with p^h elements and with $(p, m) = 1$. Recall that $\chi_{\mathfrak{p}}$ is the character of $\mathbb{F}_{p^h}^{\times}$, where $\chi_{\mathfrak{p}}(z)$ is the unique m-root of the unity satisfying $\chi_{\mathfrak{p}}(z) = z^{p^h-1/m} \bmod \mathfrak{p}$. We consider the Jacobi sum

$$J_{(a)}(\mathfrak{p}) := (-1)^{d+1} \sum_{\substack{z_1+\cdots+z_d=-1 \quad \bmod \mathfrak{p} \\ z_1,\cdots,z_d \quad \bmod \mathfrak{p}}} \chi_{\mathfrak{p}}^{a_1}(z_1) \cdots \chi_{\mathfrak{p}}^{a_d}(z_d),$$

with $a := (a_1, \cdots, a_d)$.

The map $\mathfrak{p} \to J_{(a)}(\mathfrak{p})$ defines a Hecke character for the cyclotomic field K and the ideal generated by $J_{(a)}(\mathfrak{p})$ within $\mathbb{Z}[\epsilon_m]$ is $\mathfrak{p}^{\theta(a)}$, where

$$\theta(a) := \sum_{\substack{(t,m)=1 \\ t \bmod m}} [\sum_{i=1}^{d} < \frac{t \cdot a_i}{m} >] \sigma_{-t}^{-1},$$

σ_{-t} is the automorphism of K given by $\sigma_{-t}(\epsilon_m) = \epsilon_m^{-t}$. Moreover, $|J_{(a)}(\mathfrak{p})|^2 = p^{h(s-2)}$, s being the integers $a_1, \cdots, a_d, a_1 + \cdots + a_d \neq 0 \pmod{m}$.

Here, by using previous notation $D(f(x)) = a_0 \cdot x_0 + a_1 \cdot x_1 + \cdots + a_d \cdot x_d + m \cdot D$, $a := (a_1, \cdots, a_d)$ with $1 \leq |a_i| < m$, $a_i \neq 0 \bmod \mathrm{m}$ and $a_1 + \cdots + a_d \neq 0 \bmod \mathrm{m}$. Therefore, here $s = d + 1$.

Now, from Lemma 4.3 we deduce, ([D] 5 and Proposition 10.12.1.)

Theorem 1. *The term constant of* $P_{F_{\mathfrak{p}}}^{\chi_f}(t)$ *is*

$$(-1)^{d+1} \chi_f(\sigma_M) \cdot q^g \cdot J_{(a)}(\mathfrak{p}).$$

Proof. Bearing in mind the paragraph above Remark 2.1,

$$\sigma_{z^{-1} \cdot M} = \sigma_{\eta(1, z_1^{-1}, \cdots, z_d^{-1})} \cdot \sigma_M.$$

Therefore,

$$\chi_f(\sigma_{z^{-1} \cdot M}) = \chi_{\mathfrak{p}}^{a_1}(z_1) \cdots \chi_{\mathfrak{p}}^{a_d}(z_d) \cdot \chi_f(\sigma_M),$$

where $(1, z_1, \cdots, z_d) \in k(\overline{x}_0) \times \cdots \times k(\overline{x}_d))^\times / k(\mathfrak{p})^\times$ and $[M] \in G_\mathfrak{m}$ is defined above Lemma 4.3. Note that we denote $z^{-1} \cdot M$ instead of $\eta(z^{-1}) \cdot [M]$.

By Lemma 4.3, we have that for $\mu \in \mathbb{F}_\mathfrak{p}^\times$

$$\sum_{z \in \mathcal{O}_{\mathfrak{m}'}^\times,\, \omega(z) = \mu} \chi_f(\sigma_{z^{-1} \cdot M}) =$$

$$= \chi_\mathfrak{p}^{a_1 + \cdots + a_d}(\mu) \cdot \sum_{\substack{z_1 + \cdots + z_d = -1 \mod \mathfrak{p} \\ z_1, \cdots, z_d \mod \mathfrak{p}}} \chi_\mathfrak{p}^{a_1}(z_1) \cdots \chi_\mathfrak{p}^{a_d}(z_d) \cdot \chi_f(\sigma_M).$$

The sum over $\mu \in \mathbb{F}_\mathfrak{p}^\times$ of the above terms is 0, because $a_1 + \cdots + a_d \neq 0 \mod \mathrm{m}$. Thus, by Lemma 4.3 we have:

$$\sum_{[N] \in G_\mathfrak{m}} L_\mathfrak{m}(N, 2g + d - 1) \cdot \chi_f(\sigma_N) =$$

$$= \sum_{z \in \mathcal{O}_{\mathfrak{m}'}^\times,\, \omega(z) = 1} q^g \cdot \sigma_{z^{-1} \cdot M} + \sum_{\mu \in \mathbb{F}_\mathfrak{p}^\times} q^{g-1} \sum_{z \in \mathcal{O}_{\mathfrak{m}'}^\times,\, \omega(z) = \mu} \chi_f(\sigma_{z^{-1} \cdot M}) +$$

$$+ \sum_{N + (2g + d - 1) D_1 \nsim \kappa + E} q^{g-1} \cdot \chi_f(\sigma_N).$$

Since the set of classes of level structures for the line bundle associated with M is given by $\{\eta(z^{-1}) \cdot [M]\}_{z \in \mathcal{O}_{\mathfrak{m}'}^\times} \subset G_\mathfrak{m}$, the last sum is equal to

$$\sum_{z \in \mathcal{O}_{\mathfrak{m}'}^\times,\, \omega(z) = 1} q^g \cdot \chi_f(\sigma_{z^{-1} \cdot M}) + \sum_{[N] \in G_\mathfrak{m}} q^{g-1} \cdot \chi_f(\sigma_N) = \sum_{z \in \mathcal{O}_{\mathfrak{m}'}^\times,\, \omega(z) = 1} q^g \cdot \chi_f(\sigma_{z^{-1} \cdot M})$$

and we conclude. Recall that, $\omega(z_1, \cdots, z_d) = -z_1 - \cdots - z_d$. □

Remark 5.1. As noted at the end of section 3, by [T], 3.5 we have

$$det(F_\mathfrak{p}) = \sum_{[N] \in G_\mathfrak{m}} L_\mathfrak{m}(N, 2g + d - 1) \cdot \chi_f(\sigma_N).$$

Thus, by [W] and Theorem 1

$$\mathfrak{p} \to (-1)^{d+1} \chi_f(\sigma_M)^{-1} det(F_\mathfrak{p}) = q^g \cdot J_{(a)}(\mathfrak{p}),$$

with $\mathfrak{p} \in Spec(A)$, gives a Hecke character for K.

Moreover, we also have the equality

$$\sum_{[N]\in G_{\mathfrak{m}}} L_{\mathfrak{m}}(N, 2g+d-1)\cdot\chi_f(\sigma_N) = \sum_{D,\, deg(D)=2g+d-1} \chi_{\mathfrak{p}}(f(D)),$$

where D denotes effective divisors on $X_{\mathfrak{p}}$ with support outside T. Therefore, we deduce the formula

$$\sum_{D,\, deg(D)=2g+d-1} \chi_{\mathfrak{p}}(f(D)) = (-1)^{d+1}\chi_f(\sigma_M)\cdot p^g\cdot J_{(a)}(\mathfrak{p}).$$

We have defined $f(D)$ at the end of the section 3.

For example, let X be a proper, smooth model for the elliptic curve $y^2 - x\cdot(x-1)\cdot(x-\lambda)$, defined over $A := \mathbb{Z}[\epsilon_m, \frac{1}{2m}]$, $m \neq 3$. We have that $D(y) = -3\cdot\infty + (0,0) + (1,0) + (\lambda, 0)$. Following the notation used in the above formula

$$\sum_{D,\, deg(D)=4} \chi_{\mathfrak{p}}(y(D)) = \chi_y(\sigma_M)\cdot p\cdot J_{(a)}(\mathfrak{p}).$$

D being the effective divisors on the elliptic curve, $X_{\mathfrak{p}}$, with support outside $T := \{\infty, (0,0), (1,0), (\lambda,0)\}$ and $a = (-3,1,1,1)$.

6. The p-rank of jacobians

In this section we consider m as a prime integer.

Definition 6.1. *A proper, smooth and geometrically irreducible curve Z, over a finite field $\mathbb{F}_q (q = p^h)$ has p-rank* 0 *when, for the morphism* $[p] : \mathrm{Pic}^0_{\mathrm{Z}} \to \mathrm{Pic}^0_{\mathrm{Z}}$, $\mathrm{Ker[p]_{red}} = \mathrm{Spec}(\mathbb{F}_q)$ *or, equivalently, when* $[p]$ *is purely inseparable.*

In the following proposition, we consider F_q, the q-Frobenius morphism, as a $\mathbb{Q}_l$-linear application over the $\mathbb{Q}_l$-vector space $H^1(Z, \mathbb{Q}_l)^\vee$.

Proposition 6.2. *We have that Z has p-rank* 0 *if and only if the characteristic polynomial of F_q, $t^{2\pi} + a_1 t^{2\pi-1} + \cdots + a_{2\pi-1}t + a_{2\pi} \in \mathbb{Z}[t]$, satisfies $p|a_1, \cdots, p|a_{2\pi}$. Here, π denotes the genus of Z.*

Proof. If $p|a_1, \cdots, p|a_{2\pi}$ then by considering F_q as an endomorphism of $\mathrm{Pic}^0_{\mathrm{Z}}$, we have that $F_q^{2\pi} = [p] \cdot \Phi$, where Φ is also an endomorphism of $\mathrm{Pic}^0_{\mathrm{Z}}$. Thus $[p]$ is purely inseparable because F_q is purely inseparable.

Conversely, since $\mathrm{Ker[p]_{red}} = \mathrm{Spec(F_q)}$ there exists $l \in \mathbb{N}$ such that $F_q^l(\mathrm{Ker[p]}) = \mathrm{Spec}(\mathbb{F}_\mathrm{q})$. In this way, $F_q^l = [p] \cdot \Phi$ with Φ an isogeny of $\mathrm{Pic}^0_{\mathrm{Z}}$. Thus, the characteristic polynomial of F_q is $t^{2\pi} \,\mathrm{mod}\, \mathrm{p}$. □

Let $W \to S$ be a ramified abelian covering, of group $G := \mathbb{Z}/m$, of proper, smooth and geometrically irreducible curves over $\mathbb{F}_q$, with $(m, p) = 1$. We consider, by convenience in the notation, that l and m are integer primes and that l is inert in $\mathbb{Z}[\epsilon_m]$. We consider the $\mathbb{Q}$-algebra decomposition $\mathbb{Q}[G] = \mathbb{Q} \times \mathbb{Q}(\epsilon_m)$. Let $e_1 \in \mathbb{Q}[G]$ be the idempotent element, such that $e_1 \cdot \mathbb{Q}[G] = 0 \times \mathbb{Q}[\epsilon_m]$. We consider the $\mathbb{Q}_l[G]$-module, $H^1(W, \mathbb{Q}_l)^\vee$. We denote by $p_1(t) \in \mathbb{Z}[\epsilon_m][t]$ the characteristic polynomial of F_q as a $\mathbb{Q}_l(\epsilon_m)$-endomorphism over the $\mathbb{Q}_l(\epsilon_m)$-vector space $e_1 \cdot H^1(W, \mathbb{Q}_l)^\vee$, and by $p_0(t)$ the characteristic polynomial of F_q over the $\mathbb{Q}_l$-vector space $H^1(S, \mathbb{Q}_l)^\vee$.

Lemma 6.3. *We consider $m|p-1$. If W has p-rank 0 then $p_1(0) = u \cdot p^n$, with $u \in \mathbb{Z}[\epsilon_m]$ and $n > 1 \in \mathbb{N}$.*

Proof. By the very proposition $F_q^l = [p^h] \cdot \Phi$, where Φ is an isogeny of the abelian variety $\mathrm{Pic}^0_{\mathrm{W}}$. The characteristic polynomial of Φ as a $\mathbb{Q}_l(\epsilon_m)$-endomorphism is $t^r + b_{r-1}t^{r-1} + \cdots + b_0 \in \mathbb{Z}[\epsilon_m]$. Note that F_q^l and $[p^h]$ are $\mathbb{Q}_l(\epsilon_m)$-endomorphisms and thus Φ. Therefore, the characteristic polynomial of F_q^l is $t^r + c_{r-1}t^{r-1} + \cdots + b_0 \cdot p^{r \cdot h} \in \mathbb{Z}[\epsilon_m]$. We thus have $p_1(0)^l = b_0 \cdot p^{r \cdot h}$. We conclude because the primary ideal decomposition of p in $\mathbb{Z}[\epsilon_m]$ is a product of different ideal primes, (note that $m|p-1$). □

We consider the notations of sections 1 and 2, $\mathfrak{p}$ being a prime ideal within $\mathbb{Z}[\epsilon_m]$ over p. Recall that $Y_\mathfrak{p} \to X_\mathfrak{p}$ is a cyclic ramified covering of Galois group $\mathbb{Z}/m$, and $\Sigma_{Y_\mathfrak{p}} = \Sigma_{X_\mathfrak{p}}(\sqrt[m]{f(x)})$ with $D(f(x)) = a_0 \cdot \overline{x}_0 + \cdots + a_d \cdot \overline{x}_d + m \cdot \overline{D}$. For $W = Y_\mathfrak{p}$ and $S = X_\mathfrak{p}$, we have $p_1(t) = P^{\chi_f}_{F_\mathfrak{p}}(t)$.

Theorem 2. *1) If there exists $t \in \{1, \cdots, m-1\}$ with*

$$[< \frac{a_1}{m} > + \cdots + < \frac{a_d}{m} >] \neq [< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >]$$

then when $m|p-1$(i.e p splits completely in $\mathbb{Z}[\epsilon_m]$), $\mathrm{Pic}^0_{\mathrm{Y}_\mathfrak{p}}$ is not supersingular.

2) If $X_\mathfrak{p} = \mathbb{P}_1$ and there exists $t \in \{1, \cdots, m-1\}$ with

$$[< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >] = 0$$

and $m|p-1$ then $Y_\mathfrak{p}$ has not p-rank 0.

Proof. 1) By Theorem 1, the ideal within $\mathbb{Z}[\epsilon_m]$ generated by the term constant is $(q^g \cdot J_{(a)}(\mathfrak{p}))$, thus by using [W] (8), (9) this ideal is $(q^g \cdot \mathfrak{p}^{\theta(a)})$ with

$$\theta(a) := \sum_{(t,m)=1} [< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >] \sigma^{-1}_{-t}.$$

Because

$$[< \frac{a_1}{m} > + \cdots + < \frac{a_d}{m} >] \neq [< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >]$$

for some $t \in \{1, \cdots, m-1\}$ and p splits completely in $\mathbb{Z}[\epsilon_m]$, we have that $(q^g \cdot \mathfrak{p}^{\theta(a)})$ is different to a product of "$\zeta\sqrt{q}$", and we conclude.

We prove 2) by the same arguments as in 1) but now using the above Lemma and the fact that $g = 0$. □

Lemma 6.4. *Bearing in mind the above notations with $(p, m) = 1$, $Y_\mathfrak{p}$ has p-rank 0 if and only if $X_\mathfrak{p}$ has p-rank 0 and $P^{\chi^j_f}_{F_\mathfrak{p}}(t) = t^r + p \cdot Q_j(t)$ where $Q_j(t) \in \mathbb{Z}[\epsilon_m][t]$ and deg$(Q_j(t)) < r$, for each j, $1 \leq j \leq m-1$.*

Proof. Because the characteristic polynomial of the p^h-Frobenius morphism, considered as a $\mathbb{Q}_l$-linear application of $H^1(\overline{Y}_\mathfrak{p}, \mathbb{Q}_l)^\vee$, is $p_0(t) \cdot$

$\prod_{(j,m)=1} P_{F_{\mathfrak{p}}}^{\chi_f^j}(t)$, we conclude by Proposition 6.2 and because the primary ideal decomposition of p in $\mathbb{Z}[\epsilon_m]$ is a product of different ideal primes because $(p, m) = 1$. □

Theorem 3. *1) Let p be a prime integer inert within $\mathbb{Z}[\epsilon_m]$. We have that $Y_{\mathfrak{p}}$ has p-rank 0 if and only if $X_{\mathfrak{p}}$ has p-rank 0 and*

$$\sum_{D,\mathrm{deg(D)=l}} \chi_{\mathfrak{p}}(f(D)) = 0 \operatorname{mod} \mathrm{p},$$

for each l, $1 \leq l \leq 2g + d - 2$. Note that $k(\mathfrak{p}) = \mathbb{F}_{p^{m-1}}$.

2) If we assume that p splits completely in $\mathbb{Z}[\epsilon_m]$, and for $g = 0$ that

$$[< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >] \neq 0$$

for each $t \in \{1, \cdots, m-1\}$, we have that $Y_{\mathfrak{p}}$ has p-rank 0 if and only if $X_{\mathfrak{p}}$ has p-rank 0 and

$$\sum_{D,\mathrm{deg(D)=l}} \chi_{\mathfrak{p}}(f(D))^j = 0 \operatorname{mod} \mathfrak{p}$$

for each j with $1 \leq j \leq m - 1$ and each l, $1 \leq l \leq 2g + d - 2$. Note that in this case $k(\mathfrak{p}) = \mathbb{F}_p$.

Here, D denotes the effective divisors D on $X_{\mathfrak{p}}$ with support outside T.

Proof. 1) From (*) section 3, we have

$$P_{F_{\mathfrak{p}}}^{\chi_f}(t) = \sum_{i=0}^{2g+d-1} \sum_{D,\mathrm{deg(D)=i}} \chi_{\mathfrak{p}}(f(D)) t^{2g+d-1-i}.$$

We conclude by using the above Lemma and the fact that the ideal generated in $\mathbb{Z}[\epsilon_m]$ by the constant term of this polynomial is given, by Theorem 1, by $(q^g \cdot J_{(a)}(\mathfrak{p}))$. Note that in the case of p being inert within $\mathbb{Z}[\epsilon_m]$, $\mathfrak{p} = p \cdot \mathbb{Z}[\epsilon_m]$ and $q^g J_{(a)}(\mathfrak{p}) = \epsilon_m^j \cdot p^u$ for some $1 < u$, $j \in \mathbb{N}$.

2) One proceed in the same way that 1) bearing in mind that for $g = 0$

$$[< \frac{t \cdot a_1}{m} > + \cdots + < \frac{t \cdot a_d}{m} >] \neq 0$$

for each t, $1 \leq t \leq m-1$. Thus, $(p^g \cdot J_{(a)}(\mathfrak{p})) = (q^g \cdot \mathfrak{p}^{\theta(a)}) = p \cdot I$ for some ideal $I \subset \mathbb{Z}[\epsilon_m]$. Note that $P^{\chi_f^j}_{F_\mathfrak{p}}(t) = \sigma_j(P^{\chi_f}_{F_\mathfrak{p}}(t))$, $p \cdot \mathbb{Z}[\epsilon_m] = \mathfrak{p}_1 \cdots \mathfrak{p}_{m-1}$, and the "$\sigma_j$" operate transitively on $\mathfrak{p}_1, \cdots, \mathfrak{p}_{m-1}$. Where the $\mathfrak{p}_i$ are the primes in $\mathbb{Z}[\epsilon_m]$ over p. $\square$

In the next Corollary by using part 1) of this Theorem, we obtain counterparts to the Deuring polynomial for genus greater than 1. We consider the curves defined over $\mathbb{Z}[\frac{1}{m}, \epsilon_m]$, $X = \mathbb{P}_1$ and Y associated with the plane curve

$$y^m = x^{a_1}(x-1)^{a_2}(x-\alpha_1)^{a_3} \cdots (x-\alpha_{d-2})^{a_d},$$

where the a_j are integers $\neq 0 \mod \mathrm{m}$, $a_1 + \cdots + a_d \neq 0 \,\mathrm{mod}\, \mathrm{m}$ and $\alpha_1, \cdots, \alpha_{d-2}$ are independent variables. We consider $\mathfrak{p} \subset \mathbb{Z}[1/m, \epsilon_m]$ a prime over p such that $k(\mathfrak{p}) = \mathbb{F}_{p^{m-1}}$ (i.e. p is inert within $\mathbb{Z}[\epsilon_m]$).

Corollary 6.5. *Let $Y_\mathfrak{p}$ be the proper, smooth model of the reduction at $\mathfrak{p}$ of Y, $k(\mathfrak{p}) = \mathbb{F}_{p^{m-1}}$. Accordingly, $Y_\mathfrak{p}$ has p-rank 0 if and only if*

$$\sum_{\substack{q(x) \\ \deg(q(x))=l}} [q(0)^{a_1} q(1)^{a_2} q(\alpha_1)^{a_3} \cdots q(\alpha_{d-2})^{a_d}]^{p^{m-1}-1/m} = 0 \,\mathrm{in}\, \mathbb{F}_{\mathrm{p}^{\mathrm{m}-1}},$$

for each l, with $1 \leq l \leq d-2$. Here, $q(x)$ denotes monic polynomials $q(x) \in \mathbb{F}_{p^{m-1}}[x]$

Proof. It suffices to consider in the above theorem $X_\mathfrak{p} = \mathbb{P}_1$ with $\Sigma_{X_\mathfrak{p}} = \mathbb{F}_{p^{m-1}}(x)$, $T := (x \cdot (x-1) \cdot (x-\alpha_1) \cdots (x-\alpha_{d-2}))_0$ and also to consider that the effective divisors of degree l on $Spec(\mathbb{F}_{p^{m-1}}[x])$ are given by the zero locus of monic polynomials $q(x)$ of degree l.

$\square$

Note that the system of equations of this corollary is a system on $\mathbb{F}_{p^{m-1}}$ with $d-2$ variables and $d-2$ equations.

For example, the hyperelliptic curve defined over $\mathbb{F}_p$

$$y^2 - x(x-1)(x-\alpha_1)(x-\alpha_2)(x-\alpha_3)$$

has p-rank 0 with $p \neq 2$, if and only if

$$\sum_{a\in\mathbb{F}_p}[a(1+a)\prod_{i=1}^{3}(\alpha_i+a)]^{p-1/2}=0$$

$$\sum_{a,b\in\mathbb{F}_p}[a(1+b+a)\prod_{i=1}^{3}(\alpha_i^2+\alpha_i\cdot b+a)]^{p-1/2}=0$$

$$\sum_{a,b,c\in\mathbb{F}_p}[a(1+c+b+a)\prod_{i=1}^{3}(\alpha_i^3+\alpha_i^2\cdot c+\alpha_i\cdot b+a)]^{p-1/2}=0.$$

The next Corollary is an application of part 2) of the above Theorem. We consider $\mathfrak{p}\subset\mathbb{Z}[1/m,\epsilon_m]$ a prime over p such that $k(\mathfrak{p})=\mathbb{F}_p$ (i.e. p splits completely within $\mathbb{Z}[\epsilon_m]$), and Y the above plane curve.

Corollary 6.6. *Let $Y_{\mathfrak{p}}$ be the proper, smooth model of the reduction at $\mathfrak{p}$ of Y, where p splits completely in $\mathbb{Z}[1/m,\epsilon_m]$ and*

$$[<\frac{t\cdot a_1}{m}>+\cdots+<\frac{t\cdot a_d}{m}>]\neq 0$$

for each t, $1\leq t\leq m-1$. We have that $Y_{\mathfrak{p}}$ has p-rank 0 if and only if

$$\sum_{\substack{q(x)\\ \deg(q(x))=l}}[q(0)^{a_1}q(1)^{a_2}q(\alpha_1)^{a_3}\cdots q(\alpha_{d-2})^{a_d}]^{j(p-1)/m}=0 \bmod \mathfrak{p},$$

for each l and j with $1\leq l\leq d-2$ and $1\leq j\leq m-1$, respectively. Now, $q(x)$ denotes monic polynomials $q(x)\in\mathbb{F}_p[x]$.

Proof. One proceeds in the same way as in the above corollary by using part 2) of the above Theorem.

□

As an example of this corollary one proves that the desingularization of the curve, defined over $\mathbb{F}_7$, $y^3-x\cdot(x-1)\cdot(x-\alpha)^2$ has 7-rank $\neq 0$, because the system of two equations on α

$$\sum_{a\in\mathbb{F}_7}a\cdot(a-1)\cdot(a-\alpha)^2=0,\quad \sum_{a\in\mathbb{F}_7}a^2\cdot(a-1)^2\cdot(a-\alpha)^4=0$$

does not have solutions.

Acknowlegment: I would like to express my gratitude to Prof. Jesús Muñoz Díaz. I would like to thank a referee for noting a mistake in Lemma 6.3 and the important reference [D].

REFERENCES

[A1] Anderson, G. "Rank one elliptic modules A-modules and A-harmonic series", Duke Mathematical Journal. **73** (1994), pp.491-542

[A2] Anderson, G. "Torsion points on Jacobians of quotients of Fermat curves and p-adic soliton theory", Invent. Math. **118** (1994), no. 3, pp.475-492

[D] Deline, P. "Les constants des equations fonctionnelles des fonctions L", Proc. Antwerpen Conference, vol. 2; Lecture Notes in Math. **349** (Springer-Verlag 1973), pp.501-597

[L] Pries, R. "A short guide to p-torsion of abelian varieties in characteristic p", Contemporary Mathematics . **463** (2006)

[N] Noboru, A. "On supersingular cyclic quotients of Fermat curves", Comment. Math. Univ. St. Pauli . **57** (2008), no. 1, pp.65-90

[NG] Ngo, B. C. Genestier, A. "Alcôves et p-rang des varits abliennes", Ann. Inst. Fourier. **52** (2002), no. 6, pp.1665-1680

[S] Serre, J.P. "Groupes algebriques et corps de classes", Hermann (1959)

[T] Tate, J. "Les Conjectures de Stark sur Les fonction L d'Artin en s=0", Birkhauser, Boston, 1984.

[U] Ulmer, D. "Jacobi sums, Fermat Jacobians, and ranks of abelian varieties over towers of function fields.", Math. Res. Lett. **14** (2007), no. 3, pp.453-467

[W] Weil, A. "Jacobi sums as "Grssencharaktere" ", Trans. Amer. Math. Soc. **73** (1952), pp. 487-495

Álvarez Vázquez, Arturo

e-mail: aalvarez@usal.es

Departamento de Matemáticas, Universidad de Salamanca, Plaza de la Merced 1-4. Salamanca (37008). Spain.